

**OPTIMMUNE- cyclosporine ointment**  
**Schering Corporation**

-----  
**Optimmune® (0.2% Cyclosporine, USP)**  
**Ophthalmic Ointment**

**PRODUCT INFORMATION**

**For ophthalmic use in dogs only.**

**Sterile**

**CAUTION**

**US Federal law restricts this drug to use by or on the order of a licensed veterinarian.**

**DESCRIPTION**

Each gram of OPTIMMUNE® Ophthalmic Ointment contains 2 mg of cyclosporine, USP; petrolatum, USP; corn oil, NF; petrolatum and lanolin alcohol. Cyclosporine (cyclosporin A), the active ingredient of OPTIMMUNE® Ophthalmic Ointment, is a cyclic undecapeptide metabolite of the fungus *Tolypocladium inflatum gams*.

**MODE OF ACTION**

When applied ophthalmically, cyclosporine is believed to act as a local immunomodulator of diseases suspected to be immune-mediated such as keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK). In the management of KCS, the mechanism by which cyclosporine causes an increase in lacrimation is poorly understood. Clinical improvement in cases of KCS is not necessarily dependent on an increase in aqueous tear production (as measured by the Schirmer Tear Test [STT]). See

**EFFICACY.**

**INDICATIONS**

OPTIMMUNE® Ophthalmic Ointment is indicated for management of chronic keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK) in dogs.

**PRECAUTIONS**

The clinical effects of OPTIMMUNE® Ophthalmic Ointment have not been determined in dogs with KCS due to the following conditions: congenital alacrima, sulfonamide usage, canine distemper virus, metabolic disease, surgical removal of the third eyelid gland, and facial nerve paralysis with loss of the palpebral reflex. Some of the underlying conditions which may lead to KCS can be either transient (eg, facial nerve trauma) or correctable with appropriate treatment. Consequently, recovery from clinical signs attributed to KCS may be observed and treatment options may need reconsideration.

When switching to cyclosporine from another therapeutic agent (eg, frequent application of an artificial tear preparation) for KCS or CSK, it should be kept in mind that clinical efficacy is not necessarily apparent immediately after initiation of OPTIMMUNE® Ophthalmic Ointment therapy. Several days to a few weeks may be required before the clinical effects of OPTIMMUNE® Ophthalmic Ointment are of sufficient magnitude such that previously initiated therapy can be safely withdrawn. Abrupt cessation of a therapeutic agent immediately upon initiation of OPTIMMUNE® Ophthalmic Ointment therapy can result in rapid clinical relapse which may be erroneously interpreted as an adverse reaction to OPTIMMUNE® Ophthalmic Ointment.

The safety of OPTIMMUNE® Ophthalmic Ointment has not been determined in cases of preexisting viral or fungal ocular infections. It is recommended that in such cases, OPTIMMUNE® Ophthalmic Ointment therapy be delayed until the fungal/viral ocular infection has been successfully treated.

The safety of OPTIMMUNE® Ophthalmic Ointment in puppies, pregnant bitches, or dogs used for breeding has not been determined.

## **EFFICACY**

1. **KCS** A well-controlled clinical field trial was conducted by veterinary ophthalmologists in 9 states and included 132 dogs afflicted with KCS of which 124 were evaluated for efficacy. Dogs were randomly assigned to BID treatment with either 0.2% (OPTIMMUNE® Ophthalmic Ointment) or 0% (placebo vehicle) cyclosporine ophthalmic ointment for 12 weeks. Treatment with OPTIMMUNE® Ophthalmic Ointment resulted in an average 8 to 9 mm increase in STT by the end of the study period (vs 3 to 4 mm for the placebo vehicle). Most of the increase in STT, approximately 6 mm, occurred in the first week of therapy. Some dogs improved clinically (ie, exhibited a decrease in conjunctival and/or corneal pathology) without an increase in STT values. This is thought to occur through suppression of inflammation by cyclosporine on the ocular surface. In this clinical field trial, OPTIMMUNE® Ophthalmic Ointment therapy was also associated with an improvement in clinical signs in comparison to the placebo. Blepharitis, blepharospasm, and "other signs of ocular discomfort" (eg, pawing at eyes), were markedly reduced. Improvement in conjunctival health as manifested by reduced conjunctival hypertrophy, reduced hyperemia, reduced conjunctival discharge volume, and improved character of discharge was evident. Improvement in corneal health as manifested by improved corneal surface contour, reduced corneal edema and corneal neovascularization was also noted. Overall improvement was noted in 81% of eyes treated with OPTIMMUNE® Ophthalmic Ointment.

Withdrawal of OPTIMMUNE® Ophthalmic Ointment therapy resulted in rapid clinical regression in all but one test eye indicating the need for long-term continual therapy for almost all cases of chronic KCS.

2. **CSK** The efficacy of OPTIMMUNE® Ophthalmic Ointment was determined in a historical controlled clinical field trial conducted by veterinary ophthalmologists in four countries and included 36 dogs afflicted with CSK. Dogs, primarily German shepherds, a breed disposed to CSK (German shepherd pannus), were treated twice daily with OPTIMMUNE® Ophthalmic Ointment for 6 weeks. Clinical improvement was noted by the investigators in 90.3% of eyes treated with OPTIMMUNE® Ophthalmic Ointment when compared to baseline.

## **SAFETY**

A target animal safety study and clinical field studies with OPTIMMUNE® Ophthalmic Ointment showed a wide safety margin in adult dogs. In the 6-month target animal safety study, dogs were subjected twice daily to up to 10 times the approved concentration of OPTIMMUNE® Ophthalmic Ointment. No apparent toxicity or adverse reactions were observed. Dogs in this study were vaccinated with commercially available vaccines. No effect on antibody titer response was noted. Epiphora was noted in all groups, including the placebo group, and was not associated with any inflammatory change, nor was there any correlation to gross and histopathological changes.


## **ADVERSE REACTIONS**

In the KCS clinical field trial, there were 20 adverse reactions reported out of 132 cases enrolled. This corresponds to an adverse reaction rate of 12.9% (13 of 101 cases) for OPTIMMUNE® Ophthalmic Ointment treated dogs and 22.6% (7 of 31) for placebo treated dogs. The reactions described were primarily ocular and periocular inflammatory reactions. These were likely a function of therapy being unable to fully control the keratoconjunctivitis, rather than a true "adverse reaction." Similarly, in the CSK trial, of 36 cases evaluated for safety, adverse reactions were noted in 2 animals (5.6%). One

involved transient hyperemia, epiphora, and mild discomfort of the eye. The other involved periocular/palpebral inflammation and mild alopecia.

On rare occasion, instillation of OPTIMMUNE® Ophthalmic Ointment may be associated with local irritation as manifested by periocular redness, lid spasm, and excessive rubbing. As the eyes of dogs with KCS often demonstrate considerable inflammation, it will be difficult to determine whether this local irritation constitutes a hypersensitivity to OPTIMMUNE® Ophthalmic Ointment. If this ocular irritation persists beyond 7 days, hypersensitivity to a component of OPTIMMUNE® Ophthalmic Ointment should be suspected and therapeutic options reassessed.

## DOSAGE AND ADMINISTRATION

Remove debris with suitable nonirritating solutions. Apply a 1/4 inch strip of  ointment to the affected eye(s) every 12 hours. The ointment may be placed directly on the cornea or into the conjunctival sac.

It is recommended that dogs exhibiting chronic recurring conjunctivitis be tested for adequate tear production to determine if they are suffering from early stages of chronic KCS.

For best results in treating KCS, cyclosporine ophthalmic ointment should be administered early in the course of the disease before irreversible damage to the lacrimal tissue, or dense corneal scarring or pigmentation occurs.

Dogs afflicted with KCS or CSK will most likely require lifelong consistent therapy (see **EFFICACY** section above). For CSK, because environmental factors such as ultraviolet (UV) radiation are implicated in the pathogenesis, clinical signs may subside in the winter months when light intensity is reduced or if the dog is moved to a lower altitude, or indoors, and thus exposed to less UV radiation.<sup>1</sup>

In cases refractory to cyclosporine, the diagnosis should be reevaluated and a different course of therapy considered. Periodic reassessment of the need for OPTIMMUNE® Ophthalmic Ointment therapy is recommended.

## HOW SUPPLIED

OPTIMMUNE® Ophthalmic Ointment is available in a 3.5 g tube, carton of 6 (NDC 0061-1088-01).

## STORAGE CONDITIONS

Store between 2° and 25° C (36° and 77° F).

**KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.**

## REFERENCE

<sup>1</sup>Roberts, Steven M. *Pannus*. In: Kirk's Current Veterinary Therapy XII, Small Animal Practice. Philadelphia: W.B. Saunders Co; 1995:1245-1248.

Made in France

Optimmune is the property of Intervet International B.V. or affiliated companies or licensors and is protected by copyrights, trademark and other intellectual property laws.

Copyright © 2011 Intervet International B.V. All rights reserved.

Rev. 4/11

114909 R6

NADA # 141-052, Approved by FDA

**PRINCIPAL DISPLAY PANEL - 3.5 g Tube Carton**

NDC 0061-1088-01

**Optimmune®  
(0.2% Cyclosporine, USP)  
Ophthalmic Ointment**

NADA #141-052, Approved by FDA.

**Keep Out of Reach of Children**

*SP Intervet  
Schering-Plough Animal Health*

**6 Tubes  
3.5 g Each  
Sterile**

**Caution:** Federal law restricts  
this drug to use by or on the  
order of a licensed veterinarian.

NDC 0061-1088-01

**Optimmune<sup>®</sup>**  
(0.2% Cyclosporine, USP)  
Ophthalmic Ointment

NADA #141-052, Approved by FDA.

**Keep Out of Reach of Children**

 **Intervet**  
Schering-Plough Animal Health

**Caution:** Federal law restricts  
this drug to use by or on the  
order of a licensed veterinarian.

**6 Tubes**  
**3.5 g Each**  
**Sterile**

**Optimmune.**  
(0.2% Cyclosporine, USP)  
Ophthalmic Ointment

**For ophthalmic use in dogs only.**

Optimmune is the property of Intervet  
International B.V. or affiliated companies or  
licensors and is protected by copyrights,  
trademark and other intellectual property laws.  
Copyright © 2011 Intervet International B.V.  
All rights reserved.



Lot MBKA  
Exp  
Ophthalmic Ointment  
(0.2% Cyclosporine, USP)  
**Optimmune®**  
For ophthalmic use in dogs only.  
6 Tubes  
3.5 g Each  
Sterile

**For ophthalmic use in dogs only.**

**Indications:** OPTIMMUNE® (0.2% cyclosporine, USP) Ophthalmic Ointment is indicated for management of chronic keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK) in dogs.

**Each gram of ointment contains:** 2mg of cyclosporine, USP; petrolatum, USP; com oil, NF; petrolatum and lanolin alcohol.

**Dosage and Administration:** The eye should be properly cleansed of debris with suitable nonirritating solutions.

Apply approximately 1/4-inch strip of ointment to the affected eye(s) every 12 hours.

**Read accompanying directions carefully.  
Store between 2° and 25°C (36° and 77°F).**

**Optimmune.**  
(0.2% Cyclosporine, USP)  
Ophthalmic Ointment

**For ophthalmic use in dogs only.**



8 713184 107886 >

11 4545 R2  
SEG-CA35Gx6\_01\_R4  
84x47x100mm  
PMS 485C black

**OPTIMMUNE**  
cyclosporine ointment

<b>Product Information</b>				
<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:0061-1088	
<b>Route of Administration</b>	OPHTHALMIC			
<b>Active Ingredient/Active Moiety</b>				
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
	Cyclosporine (UNII: 83HN0GTJ6D) (Cyclosporine - UNII:83HN0GTJ6D)	Cyclosporine	2 mg in 1 g	
<b>Inactive Ingredients</b>				
	<b>Ingredient Name</b>		<b>Strength</b>	
	Paraffin (UNII: I9O0E3H2ZE)			
	Corn Oil (UNII: 8470G57WFM)			
	Petrolatum (UNII: 4T6HI2BN9U)			
	Lanolin Alcohols (UNII: 884C3FA9HE)			
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0061-1088-01	1 in 1 CARTON		
1		3.5 g in 1 TUBE		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
NADA	NADA141052	03/10/2011		

**Labeler** - Schering Corporation (001317601)

### Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Schering-Plough Sante Animale		275254357	MANUFACTURE

Revised: 6/2011

Schering Corporation