



R-Pen[®]

(penicillin G potassium)

DESCRIPTION

R-Pen[®] (penicillin G potassium) is a water-soluble antibiotic for use in drinking water for turkeys.

PACKAGING

Available in two convenient sizes:

- 25 x 0.500 billion unit packs/pail
- 12 x 1.0 billion unit packs/pail

INDICATIONS

Turkeys: For treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).



DOSAGE AND ADMINISTRATION

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days. Combine contents of entire package and approximately 1.5 pints (3 pints for 1.0 BU pack) of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

MIXING DIRECTIONS

Automatic Watering Systems: Pour the concentrate solution into a glass or plastic container, then add enough water to make 2.6 gallons (9.9 L) – 0.500 BU pack and 5.2 gallons (19.7 L) – 1.0 BU pack of solution. [This amount of solution will medicate 333 gallons (1260 L) – 0.500 BU pack and 666 gallons (2521 L) – 1.0 BU pack of drinking water.] The automatic waterer should be adjusted to deliver 1 oz (30 mL) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.

Gravity Flow Watering Systems: Pour the concentrated solution into enough water to make 333 gallons (1260 L) – 0.500 BU pack and 666 gallons (2521 L) – 1.0 BU pack of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

PRECAUTIONS

- Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.
- For best results, the treatment should be started at the first sign of infection. If improvement is not observed after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian.
- For oral use in turkeys only.

RESIDUE WARNINGS

- Treated turkeys must not be slaughtered for food during treatment and for one day after the last treatment has been completed.
- Do not use in turkeys producing eggs for human consumption.

STORAGE

Store below 86°F in a cool dry place.
Avoid excessive heat (104°F or 40°C).



R-Pen is a registered trademark of Huvepharma.

HVP-RPN-PDS-03.18



R-Pen[®]

(penicillin G potassium)

ANTIBIOTIC for Drinking Water. For Oral Use in Turkeys Only

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

Nonsterile

Mix 1 packet to:

- **R-Pen 0.500 bu:** 333 gallons
- **R-Pen 1.0 bu:** 666 gallons

Active Ingredient:

- **R-Pen 0.500 bu:** 0.500 billion units penicillin G potassium
- **R-Pen 1.0 bu:** 1.0 billion units penicillin G potassium

Gravity Flow Watering Systems:

Pour the concentrated solution into enough water to make:

- **R-Pen 0.500 bu:** 333 gallons (1260 liters)
- **R-Pen 1.0 bu:** 666 gallons (2521 liters)

of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

RESIDUE WARNINGS:

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS:

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all other medications out of the reach of children.

INDICATIONS:

For treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

DOSAGE AND ADMINISTRATION:

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

DIRECTIONS:

Combine contents of entire package with:

- **R-Pen 0.500 bu:** 1.5 pints
- **R-Pen 1.0 bu:** 3 pints

of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

Automatic Watering Systems: Pour the concentrated solution into a glass or plastic container, then

add enough water to make:

- **R-Pen 0.500 bu:** 2.6 gallons (9.9 liters) of solution. [This amount of solution will medicate 333 gallons (1260 liters) of drinking water.]
- **R-Pen 1.0 bu:** 5.2 gallons (19.7 liters) of solution. [This amount of solution will medicate 666 gallons (2521 liters) of drinking water.]

The automatic waterer should be adjusted to deliver 1 oz (30 ml) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.

Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

Restricted Drug (California).
Use only as directed.

Not for human use.

Keep Out of Reach of Children
ANADA 200-106, Approved by FDA