Vetprofen Flavored Tat **DO NOT REMOVE** INFORMATION **DOG OWNER**



Dog Owner Information about **Vetprofen Flavored Tab** (Carprofen Flavored Tablets)

for Osteoarthritis and Post-Surgical Pain

This summary contains important information about Vetprofen Flavored Tab. You should read this information before you start giving your dog Vetprofen Flavored Tab and review it each time the prescription is refilled. This sheet is provided only as a summary and does not take the place of instructions from your veterinarian. Talk to your veterinarian if you do not understand any of this information or if you want to know more about Vetprofen Flavored Tab.

What is Vetprofen Flavored Tab?

Vetprofen Flavored Tab is a nonsteroidal anti-inflammatory drug (NSAID) that is used to reduce pain and inflammation (soreness) due to osteoarthritis and pain following surgery in dogs. Vetprofen Flavored Tab is a prescription drug for dogs. It is available as a tablet and is given to dogs by mouth.

Osteoarthritis (OA) is a painful condition caused by "wear and tear" of cartilage and other parts of the joints that may result in the following changes or signs in your dog:

- · Limping or lameness
- · Decreased activity or exercise (reluctance to stand, climb stairs, jump or run, or difficulty in performing these activities)
- Stiffness or decreased movement of joints

To control surgical pain (e.g. for surgeries such as spays, ear procedures or orthopedic repairs) your veterinarian may administer Vetprofen Flavored Tab before the procedure and recommend that your dog be treated for several days after going home.

What kind of results can I expect when my dog is on Vetprofen Flavored

While Vetprofen Flavored Tab is not a cure for osteoarthritis, it can relieve the pain and inflammation of OA and improve your dog's mobility.

- Response varies from dog to dog but can be quite dramatic.
- . In most dogs, improvement can be seen in a matter of days.
- If Vetprofen Flavored Tab is discontinued or not given as directed, your dog's pain and inflammation may come back.

Who should not take Vetprofen Flavored Tab?

Your dog should not be given Vetprofen Flavored Tab if he/she:

- · Has had an allergic reaction to carprofen, the active ingredient of Vetprofen Flavored Tab.
- Has had an allergic reaction to aspirin or other NSAIDs (for example deracoxib, etodolac, firocoxib, meloxicam, phenylbutazone or tepoxalin) such as hives, facial swelling, or red or itchy skin.

Vetprofen Flavored Tab should be given to dogs only.

Cats should not be given Vetprofen Flavored Tab. Call your veterinarian immediately if your cat receives Vetprofen Flavored Tab. People should not take Vetprofen Flavored Tab. Keep Vetprofen Flavored Tab and all medicines out of reach of children. Call your physician immediately if you accidentally take Vetprofen Flavored Tab.

How to give Vetprofen Flavored Tab to your dog.

Vetprofen Flavored Tab should be given according to your veterinarian's instructions. Your veterinarian will tell you what amount of Vetprofen Flavored Tab is right for your dog and for how long it should be given. Vetprofen Flavored Tab should be given by mouth and may be given with or without

What to tell/ask your veterinarian before giving Vetprofen Flavored Tab.

Talk to your veterinarian about:

- The signs of OA you have observed (for example limping, stiffness).
- · The importance of weight control and exercise in the management of
- What tests might be done before Vetprofen Flavored Tab is prescribed.
- How often your dog may need to be examined by your veterinarian.
- · The risks and benefits of using Vetprofen Flavored Tab.

Tell your veterinarian if your dog has ever had the following medical problems:

- Experienced side effects from Vetprofen Flavored Tab or other NSAIDs, such as aspirin
- Digestive upset (vomiting and/or diarrhea)
- · Liver disease
- Kidney disease
- A bleeding disorder (for example, Von Willebrand's disease)

Tell your veterinarian about:

- Any other medical problems or allergies that your dog has now or has
- · All medicines that you are giving your dog or plan to give your dog, including those you can get without a prescription.

Tell your veterinarian if your dog is:

· Pregnant, nursing or if you plan to breed your dog.

What are the possible side effects that may occur in my dog during Vetprofen Flavored Tab therapy?

Vetprofen Flavored Tab, like other drugs, may cause some side effects. Serious but rare side effects have been reported in dogs taking NSAIDs, including carprofen. Serious side effects can occur with or without warning and in rare situations result in death.

The most common NSAID-related side effects generally involve the stomach (such as bleeding ulcers), and liver or kidney problems. Look for the following side effects that can indicate your dog may be having a problem with Vetprofen Flavored Tab or may have another medical problem:

- · Decrease or increase in appetite
- Vomiting
- Change in bowel movements (such as diarrhea, or black, tarry or bloody
- Change in behavior (such as decreased or increased activity level, incoordination, seizure or aggression)
- Yellowing of gums, skin, or whites of the eyes (jaundice)
- Change in drinking habits (frequency, amount consumed)
- · Change in urination habits (frequency, color, or smell)

Change in skin (redness, scabs, or scratching)

It is important to stop therapy and contact your veterinarian immediately if you think your dog has a medical problem or side effect from Vetprofen Flavored Tab therapy. If you have additional questions about possible side effects, talk to your veterinarian.

Can Vetprofen Flavored Tab be given with other medicines?

Vetprofen Flavored Tab should not be given with other NSAIDs (for example, aspirin, deracoxib, etodolac, firocoxib, meloxicam, tepoxalin) or steroids (for example, cortisone, dexamethasone, prednisone, triamcinolone).

Tell your veterinarian about all medicines you have given your dog in the past, and any medicines that you are planning to give with Vetprofen Flavored Tab. This should include other medicines that you can get without a prescription. Your veterinarian may want to check that all of your dog's medicines can be given together.

What do I do in case my dog eats more than the prescribed amount of Vetprofen Flavored Tab?

Contact your veterinarian immediately if your dog eats more than the prescribed amount of Vetprofen Flavored Tab.

What else should I know about Vetprofen Flavored Tab?

This sheet provides a summary of information about Vetprofen Flavored Tab. If you have any questions or concerns about Vetprofen Flavored Tab, or osteoarthritis, or postoperative pain, talk to your veterinarian.

As with all prescribed medicines, Vetprofen Flavored Tab should only be given to the dog for which it was prescribed. It should be given to your dog only for the condition for which it was prescribed. It is important to periodically discuss your dog's response to Vetprofen

Flavored Tab at regular check ups. Your veterinarian will best determine if your dog is responding as expected and if your dog should continue receiving Vetprofen Flavored Tab.

To report a suspected adverse reaction call 1-800-835-9496

Issued December 2014

Manufactured by: Belcher Pharmaceuticals, LLC 12393 Belcher Road, Suite 420 Largo, Florida 33773

Distributed by: Vétoquinol U.S.A., Inc. 4250 N. Sylvania Ave. Ft. Worth, TX (USA) 76137



Vetprofen Flavored Tab

(Carprofen Flavored Tablets)

Non-steroidal anti-inflammatory drug

For oral use in dogs only

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION: Vetprofen Flavored Tab is a non-steroidal anti-inflammatory drug (NSAID) of the propionic acid class that includes ibuprofen, naproxen, and ketoprofen. Vetprofen Flavored Tab is the nonproprietary designation for a substituted carbazole, 6-chloro-ca-methyl-9H-carbazole-2-acetic acid. The empirical formula is $C_{18}H_{12}CINO_2$ and the molecular weight 273.72. The chemical structure of carprofen is:

Carprofen is a white, crystalline compound. It is freely soluble in ethanol, but practically insoluble in water

CLINICAL PHARMACOLOGY: Carprofen is a non-narcotic, non-steroidal anti-inflammatory agent with characteristic analgesic and antipyretic activity approximately equipotent to indomethacin in animal models.¹

The mechanism of action of carprofen, like that of other NSAIDs, is believed to be associated with the inhibition of cyclooxygenase activity. Two unique cyclooxygenases have been described in mammals. The constitutive cyclooxygenase, COX-1, synthesizes prostaglandins necessary for normal gastrointestinal and renal function. The inducible cyclooxygenase, COX-2, generates prostaglandins involved in inflammation.

Inhibition of COX-1 is thought to be associated with gastrointestinal and renal toxicity while inhibition of COX-2 provides anti-inflammatory activity. The specificity of a particular NSAID for COX-2 versus COX-1 may vary from species to species.³ In an in vitro study using canine cell cultures, carprofen demonstrated selective inhibition of COX-2 versus COX-1.4 Clinical relevance of these data has not been shown.

Carprofen has also been shown to inhibit the release of several prostaglandins in two inflammatory cell systems: rat polymorphonuclear leukocytes (PMN) and human rheumatoid synovial cells, indicating inhibition of acute (PMN system) and chronic (synovial cell system) inflammatory reactions.1

Several studies have demonstrated that carprofen has modulatory effects on both humoral and cellular immune responses. § 9 Data also indicate that carprofen inhibits the production of osteoclast-activating factor (OAF), PGE, and PGE₂ by its inhibitory effects on prostaglandin biosynthesis.¹

Based upon comparison with data obtained from intravenous administration, carprofen is rapidly and nearly completely absorbed (more than 90% bloavailable) when administered orally. To Peak blood plasma concentrations are achieved in 1-3 hours after oral administration of 1, 5, and 25 mg/kg to dogs. The mean terminal half-life of carprofen is approximately 8 hours (range 4.5-9.8 hours) after single oral doses varying from 1-35 mg/kg of body weight. After a 100 mg single intravenous bolus dose, the mean elimination half-life was approximately 11.7 hours in the dog. Vetprofen Flavored Tab is more than 99% bound to plasma protein and exhibits a very small volume of distribution.

Carprofen is eliminated in the dog primarily by biotransformation in the liver followed by rapid excretion of the resulting metabolites (the ester glucuronide of carprofen and the ether glucuronides of 2 phenolic metabolites, 7-hydroxy carprofen and 8-hydroxy carprofen) in the feces (70-80%) and urine (10-20%). Some enterohepatic circulation of the drug is observed.

INDICATIONS: Vetprofen Flavored Tab is indicated for the relief of pain and inflammation associated with osteoarthritis and for the control of postoperative pain associated with soft tissue and orthopedic surgeries

CONTRAINDICATIONS: Vetprofen Flavored Tab should not be used in dogs exhibiting previous hypersensitivity

WARNINGS: Keep out of reach of children. Not for human use. Consult a physician in cases of accidental ingestion by humans. For use in dogs only. Do not use in cats.

All dogs should undergo a thorough history and physical examination before initiation of NSAID therapy. Appropriate laboratory tests to establish hematological and serum biochemical baseline data prior to, and periodically during, administration of any NSAID should be considered. **Owners should be advised to observe** for signs of potential drug toxicity (see Information for Dog Owners, Adverse Reactions, Animal Safety and Post-Approval Experience).

PRECAUTIONS: As a class, cyclooxygenase inhibitory NSAIDs may be associated with gastrointestinal renal and hepatic toxicity. Effects may result from decreased prostaglandin production and inhibition of the enzyme cyclooxygenase which is responsible for the formation of prostaglandins from arachidonic acid. ¹¹ ¹⁴ When NSAIDs inhibit prostaglandins that cause inflammation they may also inhibit those prostaglandins which maintain normal homeostatic function. These anti-prostaglandin effects may result in clinically significant disease in patients with underlying or pre-existing disease more often than in healthy patients. ^{12,14} NSAID therapy could unmask occult disease which has previously been undiagnosed due to the absence of apparent clinical signs. Patients with underlying renal disease for example, may experience exacerbation or decompensation of their renal disease while on NSAID therapy.¹¹⁻¹⁴ The use of parenteral fluids during surgery should be considered to reduce the potential risk of renal complications when using NSAIDs perioperatively.

Carprofen is an NSAID, and as with others in that class, adverse reactions may occur with its use. The most frequently reported effects have been gastrointestinal signs. Events involving suspected renal, hematologic, neurologic, dermatologic, and hepatic effects have also been reported. Patients at greatest risk for renal toxicity are those that are dehydrated, on concomitant diuretic therapy, or those with renal, cardiovascular, and/or hepatic dysfunction. Concurrent administration of potentially nephrotoxic drugs should be approached cautiously, with appropriate monitoring. Concomitant use of Vetprofen Flavored Tab with other anti-inflammatory drugs, such as other NSAIDs or corticosteroids, should be avoided because of the potential increase of adverse reactions, including gastrointestinal ulcerations and/or perforations. Sensitivity to drug-associated adverse reactions varies with the individual patient. Dogs that have experienced adverse reactions from one NSAID may experience adverse reactions from another NSAID. Vetprofen Flavored Tab treatment was not associated with renal toxicity or gastrointestinal ulceration in well controlled safety studies of up to ten times the dose in healthy dogs.

Vetprofen Flavored Tab is not recommended for use in dogs with bleeding disorders (e.g., Von Willebrand's disease), as safety has not been established in dogs with these disorders. The safe use of Vetprofen Flavored Tab in animals less than 6 weeks of age, pregnant dogs, dogs used for breeding purposes, or in lactating bitches has not been established. Studies to determine the activity of Vetprofen Flavored Tab when administered concomitantly with other protein-bound or similarly metabolized drugs have not been conducted. Drug compatibility should be monitored closely in patients requiring additional therapy. Such drugs commonly used include cardiac, anticonvulsant and behavioral medications. It has been suggested that treatment with carprofen may reduce the level of inhalant anesthetics needed.¹⁵

If additional pain medication is warranted after administration of the total daily dose of Vetprofen Flavored Tab, alternative analgesia should be considered. The use of another NSAID is not recommended. Consider appropriate washout times when switching from one NSAID to another or when switching from corticosteroid

INFORMATION FOR DOG OWNERS:

Vetprofen Flavored Tab, like other drugs of its class, is not free from adverse reactions. Owners should be advised of the potential for adverse reactions and be informed of the clinical signs associated with drug intolerance. Adverse reactions may include decreased appetite, vomiting, diarrhea, dark or tarry stools, ncreased water consumption, increased urination, pale gums due to anemia, yellowing of gums, skin or white of the eve due to jaundice, lethargy, incoordination, seizure, or behavioral changes, Serious adverse reactions sociated with this drug class can occur without warning and in rare situations result in death (see Adverse Reactions). Owners should be advised to discontinue Vetprofen Flavored Tab therapy and contact their veterinarian immediately if signs of intolerance are observed. The vast majority of patients with drug related adverse reactions have recovered when the signs are recognized, the drug is withdrawn, and veterinary care, if appropriate, is initiated. Owners should be advised of the importance of periodic follow up for all dogs during administration of any NSAID.

ADVERSE REACTIONS: During investigational studies of osteoarthritis with twice daily administration of 1 mg lb. no clinically significant adverse reactions were reported. Some clinical signs were observed during field studies (n=297) which were similar for carprofen- and placebo-treated dogs. Incidences of the following were observed in both groups: vomiting (4%), diarrhea (4%), changes in appetite (3%), lethargy (1.4%), behavioral changes (1%), and constipation (0.3%). The product vehicle served as control.

There were no serious adverse events reported during clinical field studies of osteoarthritis with once daily administration of $2\,\mathrm{mg/lb}$. The following categories of abnormal health observations were reported. The product

Percentage of Dogs with Abnormal Health Observations Reported in Osteoarthritis Field Study (2 mg/lb once daily)			
Observation	Carprofen (n=129)	Placebo (n=132)	
Inappetence	1.6	1.5	
Vomiting	3.1	3.8	
Diarrhea/Soft stool	3.1	4.5	
Behavior change	0.8	0.8	
Dermatitis	0.8	0.8	
PU/PD	0.8		
SAP increase	7.8	8.3	
ALT increase	5.4	4.5	
AST increase	2.3	0.8	
BUN increase	3.1	1.5	
Bilirubinuria	16.3	12.1	
Ketonuria	14.7	9.1	

Clinical pathology parameters listed represent reports of increases from pre-treatment values; medical judgment is necessary to determine clinical relevance.

During investigational studies of surgical pain for the tablet formulation, no clinically significant adverse reactions were reported. The product vehicle served as control.

Observation*	Carprofen (n=148)	Placebo (n=149)
/omitina	10.1	13.4
Diarrhea/Soft stool	6.1	6.0
Cular disease	2.7	0
nappetence	1.4	0
ermatitis/skin lesion	2.0	1.3
lysrhythmia	0.7	0
Apnea	1.4	0
Oral/periodontal disease	1.4	0
yrexia	0.7	1.3
Irinary tract disease	1.4	1.3
Vound drainage	1.4	0

* A single dog may have experienced more than one occurrence of an event

Although not all adverse reactions are reported, the following adverse reactions are based on voluntary post-approval adverse drug experience reporting. The categories of adverse reactions are listed in decreasing order of frequency by body system.

Gastrointestinal: Vomiting, diarrhea, constipation, inappetence, melena, hematemesis, gastrointestinal ulceration, gastrointestinal bleeding, pancreatitis.

Hepatic: Inappetence, vomiting, jaundice, acute hepatic toxicity, hepatic enzyme elevation, abnormal liver function test(s), hyperbilirubinemia, bilirubinuria, hypoalbuminemia. Approximately one-fourth of hepatic reports were in Labrador Retrievers.

Neurologic: Ataxia, paresis, paralysis, seizures, vestibular signs, disorientation.

Urinary: Hematuria, polyuria, polydipsia, urinary incontinence, urinary tract infection, azotemia, acute renal failure, tubular abnormalities including acute tubular necrosis, renal tubular acidosis, glucosuria.

Behavioral: Sedation, lethargy, hyperactivity, restlessness, aggressiveness.

Hematologic: Immune-mediated hemolytic anemia, immune-mediated thrombocytopenia, blood loss anemia, epistaxis.

Dermatologic: Pruritus, increased shedding, alopecia, pyotraumatic moist dermatitis (hot spots), necrotizing panniculitis/vasculitis, ventral ecchymosis

Immunologic or hypersensitivity: Facial swelling, hives, erythema.

In rare situations, death has been associated with some of the adverse reactions listed above.

To report a suspected adverse reaction call 1-800-835-9496.

DOSAGE AND ADMINISTRATION: Always provide Client Information Sheet with prescription. Carefully consider the potential benefits and risk of Vetprofen Flavored Tab and other treatment options before deciding to use Vetprofen Flavored Tab. Use the lowest effective dose for the shortest duration consistent with individual response. The recommended dosage for oral administration to dogs is 2 mg/lb (4.4 mg/kg) of body weight daily. The total daily dose may be administered as 2 mg/lb of body weight once daily or divided and administered as 1 mg/lb (2.2 mg/kg) twice daily. For the control of postoperative pain, administer approximately 2 hours before the procedure. Tablets are scored and dosage should be calculated in half-tablet increments.

EFFECTIVENESS: Confirmation of the effectiveness of Vetprofen Flavored Tab for the relief of pain and inflammation associated with osteoarthritis, and for the control of postoperative pain associated with soft tissue and orthopedic surgeries was demonstrated in 5 placebo-controlled, masked studies examining the antiinflammatory and analgesic effectiveness of Vetprofen Flavored Tab in various breeds of dogs.

Separate placebo-controlled, masked, multicenter field studies confirmed the anti-inflammatory and analgesic separate placetor-combinet, masket, multicerien returbuscules commined the anti-initialitiation and analysis effectiveness of Vetprofen Flavored Tab when dosed at 2 mg/lb note daily or when divided and administered at 1 mg/lb twice daily. In these two field studies, dogs diagnosed with osteoarthritis showed statistically significant overall improvement based on lameness evaluations by the veterinarian and owner observations when administered Vetprofen Flavored Tab at labeled doses.

Separate placebo-controlled, masked, multicenter field studies confirmed the effectiveness of Vetprofen Separate placebor-controlled, impacked, initiatellier field solutions continue the effectiveness of verpriore Flavored Tab for the control of postoperative pain when dosed at 2 mg/lb once daily in various breeds of dogs. In these studies, dogs presented for ovariohysterectomy, cruciate repair and aural surgeries were administered Vetprofen Flavored Tab preoperatively and for a maximum of 3 days (soft tissue) or 4 days (orthopedic) postoperatively. In general, dogs administered Vetprofen Flavored Tab showed statistically significant improvement in pain scores compared to controls.

ANIMAL SAFETY: Laboratory studies in unanesthetized dogs and clinical field studies have demonstrated that Vetprofen Flavored Tab is well tolerated in dogs after oral administration.

In target animal safety studies, Vetprofen Flavored Tab was administered orally to healthy Beagle dogs at 1, 3, and 5 mg/lb twice daily (1, 3 and 5 times the recommended total daily dose) for 42 consecutive days with no significant adverse reactions. Serum albumin for a single female dog receiving 5 mg/lb twice daily decreased to 2.1 g/dL after 2 weeks of treatment, returned to the pre-treatment value (2.6 g/dL) after 4 weeks of treatment, and was 2.3 g/dL at the final 6-week evaluation. Over the 6-week treatment period, black or bloody stools were observed in 1 dog (1 incident) treated with 1 mg/lb twice daily and in 1 dog (2 incidents) treated with 3 mg/lb twice daily. Redness of the colonic mucosa was observed in 1 male that received 3 mg/lb twice daily.

Two of 8 dogs receiving 10 mg/lb orally twice daily (10 times the recommended total daily dose) for 14 days exhibited hypoalbuminemia. The mean albumin level in the dogs receiving this dose was lower (2.38 g/dL) than each of 2 placebo control groups (2.88 and 2.93 g/dL, respectively). Three incidents of black or bloody stool were observed in 1 dog. Five of 8 dogs exhibited reddened areas of duodenal mucosa on gross pathologic examination. Histologic examination of these areas revealed no evidence of ulceration, but did show minimal congestion of the lamina propria in 2 of the 5 dogs.

In separate safety studies lasting 13 and 52 weeks, respectively, dogs were administered orally up to 11.4 mg/ lb/day (5.7 times the recommended total daily dose of 2 mg/lb) of carprofen. In both studies, the drug was well tolerated clinically by all of the animals. No gross or histologic changes were seen in any of the treated animals. In both studies, dogs receiving the highest doses had average increases in serum L-alanine aminotransferase (ALT) of approximately 20 IU.

In the 52 week study, minor dermatologic changes occurred in dogs in each of the treatment groups but not in the control dogs. The changes were described as slight redness or rash and were diagnosed as nonspecific dermatitis. The possibility exists that these mild lesions were treatment related, but no dose relationship was

Clinical field studies were conducted with 549 dogs of different breeds at the recommended oral doses for 14 days (297 dogs were included in a study evaluating 1 mg/lb twice daily and 252 dogs were included in a separate study evaluating 2 mg/lb once daily). In both studies the drug was clinically well tolerated and the incidence of clinical adverse reactions for Vetprofen Flavored Tab-treated animals was no higher than placebo-treated animals (placebo contained inactive ingredients found in Vetprofen Flavored Tab). For animals receiving 1 mg/lb twice daily, the mean post-treatment serum ALT values were 11 IU greater and 9 IU less than pre-treatment values for dogs receiving Vetprofen Flavored Tab and placebo, respectively. Differences were not statistically significant. For animals receiving 2 mg/lb once daily, the mean post-treatment serum ALT values were 4.5 IU greater and 0.9 IU less than pre-treatment values for dogs receiving Vetprofen Flavored Tab and placebo, respectively. In the latter study, 3 Vetprofen Flavored Tab-treated dogs developed a 3-fold or greater increase in (ALT) and/or (AST) during the course of therapy. One placebo-treated dog had a greater than 2-fold increase in ALT. None of these animals showed clinical signs associated with laboratory value changes. Changes in the clinical laboratory values (hematology and clinical chemistry) were not considered clinically significant. The 1 mg/lb twice daily course of therapy was repeated as needed at 2-week intervals in 244 dogs, some for as long as 5 years.

Clinical field studies were conducted in 297 dogs of different breeds undergoing orthopedic or soft tissue surgery. Dogs were administered 2 mg/lb of Vetprofen Flavored Tab two hours prior to surgery then once daily, as needed for 2 days (soft tissue surgery) or 3 days (orthopedic surgery). Vetprofen Flavored Tab was well tolerated when used in conjunction with a variety of anesthetic-related drugs. The type and severity of abnormal health observations in Vetprofen Flavored Tab- and placebo-treated animals were approximately equal and few in number (see Adverse Reactions). The most frequent abnormal health observation was vomiting and was observed at approximately the same frequency in Vetprofen Flavored Tab- and placebo-treated animals. Changes in clinicopathologic indices of hematopoietic, renal, hepatic, and clotting function were not clinically significant. The mean post-treatment serum ALT values were 7.3 IU and 2.5 IU less than pre-treatment values for dogs receiving Vetprofen Flavored Tab and placebo, respectively. The mean post-treatment AST values were 3.1 IU less for dogs receiving Vetprofen Flavored Tab and 0.2 IU greater for dogs

STORAGE: Store at controlled room temperature 15°C - 30°C (59°F - 86°F).

HOW SUPPLIED: Vetprofen Flavored Tab are scored, and contain 25 mg, 75 mg, or 100 mg of carprofen per

REFERENCES:

- 1. Baruth H, et al: In Anti-Inflammatory and Anti-Rheumatic Drugs, Vol. II, Newer Anti-Inflammatory Drugs, Rainsford KD, ed. CRC Press, Boca Raton, pp. 33-47, 1986.
- 2. Vane JR, Botting RM: Mechanism of action of anti-inflammatory drugs. Scand J Rheumatol 25:102, pp. 9-21. 3. Grossman CJ, Wiseman J, Lucas FS, et al: Inhibition of constitutive and inducible cyclooxygenase activity in human platelets and mononuclear cells by NSAIDs and COX-2 inhibitors. Inflammation Research 44:253-257,
- 4. Ricketts AP, Lundy KM, Seibel SB: Evaluation of selective inhibition of canine cyclooxygenase 1 and 2 by carprofen and other nonsteroidal anti-inflammatory drugs. Am J Vet Res 59:11, pp. 1441-1446, November
- 5. Ceuppens JL, et al: Non-steroidal anti-inflammatory agents inhibit the synthesis of IgM rheumatoid factor in vitro, Lancet 1:528, 1982,
- 6. Ceuppens JL, et~al: Endogenous prostaglandin E_2 enhances polyclonal immunoglobulin production by ionically inhibiting T suppressor cell activity. *Cell Immunol* 70:41, 1982. 7. Schleimer RP, et al: The effects of prostaglandin synthesis inhibition on the immune response
- Immunopharmacology 3:205, 1981. 8. Leung KH, et al: Modulation of the development of cell mediated immunity: possible roles of the products
- of cyclooxygenase and lipoxygenase pathways of arachidonic acid metabolism. *Int J Immunopharmacology* 4:195, 1982. 9. Veit BC: Immunoregulatory activity of cultured-induced suppressor macrophages. Cell Immunol 72:14, 1982.
- 10. Schmitt M, et al: Biopharmaceutical evaluation of carprofen following single intravenous, oral, and rectal doses in dogs. Biopharm Drug Dispos 11(7): 585-94, 1990. 11. Kore AM: Toxicology of nonsteriodal anti-inflammatory drugs. Veterinary Clinics of North America, Small
- 12. Binns SH: Pathogenesis and pathophysiology of ischemic injury in cases of acute renal failure. Compend for Cont Ed 16:1. January 1994.
- 13. Boothe DM: Prostaglandins: Physiology and clinical implications. Compend for Cont Ed 6:11, November
- $14. \ Rubin \ SI: Nonsteroidal \ anti-inflammatory \ drugs, \ prostaglandins, \ and \ the \ kidney. \ \textit{JAVMA} \ 188:9, \ May \ 1986.$
- 15. Ko CH, Lange DN, Mandsager RE, et al: Effects of butorphanol and carprofen on the minimal alveolar concentration of isoflurane in dogs. *JAVMA* 217:1025-1028, 2000. For a copy of the Safety Data Sheet (SDS) or to report adverse reactions call: 1-800-835-9496.



Belcher Pharmaceuticals, LLC Vetoquinol U.S.A., Inc. 4250 N. Sylvania Ave. Ft. Worth, TX (USA) 76137 12393 Belcher Road, Suite 420



ANADA # 200-578, Approved by FDA

Manufactured by:

Largo, Florida 33773